


00000000



Drug Facts

Active ingredient (in each caplet) Purpose
Acetaminophen 650 mg..... Pain reliever

Uses
temporarily relieves minor aches and pains due to:
 ■ arthritis
 ■ the common cold
 ■ headache
 ■ toothache
 ■ muscular aches
 ■ backache
 ■ menstrual cramps


Warnings
Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain reliever/fever reducers. Acetaminophen may cause liver damage.

Do not use
 ■ with any other product containing acetaminophen

Stop use and ask a doctor if
 ■ new symptoms occur
 ■ redness or swelling is present
 ■ pain gets worse or lasts for more than 10 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdoses) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.



NEW LABEL INFORMATION

Lasts up to 8 Hours



TYLENOL[®] Pain Reliever

ARTHRITIS PAIN

Contains Acetaminophen Extended Release
FOR THE TEMPORARY RELIEF OF MINOR ARTHRITIS PAIN

100 Caplets* - 650 mg EACH
*Capsule-Shaped Tablets

Extended Relief Caplets



NEW LABEL INFORMATION

Lasts up to 8 Hours

TYLENOL[®] Pain Reliever

ARTHRITIS PAIN

Contains Acetaminophen Extended Release
FOR THE TEMPORARY RELIEF OF MINOR ARTHRITIS PAIN

100 Caplets* - 650 mg EACH
*Capsule-Shaped Tablets

Extended Relief Caplets

NDC 50580-112-101

Lasts up to 8 Hours


TYLENOL[®] Pain Reliever

ARTHRITIS PAIN

Contains Acetaminophen Extended Release
FOR THE TEMPORARY RELIEF OF MINOR ARTHRITIS PAIN

Extended Relief Caplets

McNeil
 McNeil Consumer & Specialty Pharmaceuticals
 DIVISION OF MCNEIL - PPC, INC.
 FORT WASHINGTON, PA 19034 USA
 © MCN-PPC, Inc. '02 U.S. Pat. No. 4820522
 www.tylenol.com



Drug Facts (continued)

Directions
 ■ do not take more than directed (see overdose warning)
 ■ take 2 caplets every 8 hours with water
 ■ swallow whole - do not crush, chew or dissolve
 ■ do not take more than 6 caplets in 24 hours
 ■ do not use for more than 10 days unless directed by a doctor


adults
 under 18 years of age ■ ask a doctor

Other information
 ■ do not use if carton is opened or neck wrap or foil liner seal imprinted with "Safety Seal[®]" is broken
 ■ store at 20 - 25°C (68 - 77°F)
 ■ avoid excessive heat 40°C (104°F)
 ■ see end panel for lot number and expiration date

Inactive ingredients
 corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin

Questions or comments?
 call toll-free 1-877-TYLENOL (1-877-965-9665)

What makes Tylenol[®] Arthritis Pain Extended Relief Caplets different?
 • Uses a unique, patented 2-layer caplet. The first layer dissolves quickly to provide prompt relief while the second layer is time released to provide up to 8 hours of relief.
 • For more information or questions, visit our website www.tylenol.com.
 • The makers of Tylenol[®] do not manufacture store brands.



Labeling Format Information:

Fonts: Helvetica regular, bold and bold italic.	
Drug Facts: 8.25 pt	Leading: 6.5 pt
Header: 8 pt	Bullets: 5 pt
Subheader: 6 pt	Barlines: 2.5 pt
Body Text: 6 pt	Hairlines: .5 pt
Drug Facts (continued): 8 pt	
Horizontal Scale: 75-80%	Average Kerning: 0

CTN TY ARTH ER CPLT 100 3004009 A-1
 OCTOBER 16, 2002

<p>Active ingredient (in each caplet) Acetaminophen 650 mg Purpose Pain reliever</p>	<p>READ THE LABEL DO NOT TAKE MORE THAN 6 CAPLETS IN 24 HOURS NDC 50580-112-10</p>	<p>adults as well as for children even if you do not notice any signs or symptoms.</p>
<p>Uses temporarily relieves minor aches and pains due to: ■ arthritis ■ the common cold ■ headache ■ toothache ■ muscular aches ■ backache ■ menstrual cramps</p> <p>Warnings Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take acetaminophen or other pain relievers/fever reducers. Acetaminophen may cause liver damage. Do not use ■ with any other product containing acetaminophen ■ Stop use and ask a doctor if ■ new symptoms occur ■ redness or swelling is present ■ pain gets worse or lasts for more than 10 days ■ If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for</p>	<p>DO NOT USE WITH OTHER MEDICINES CONTAINING ACETAMINOPHEN</p> <p><i>Lasts up to 8 Hours</i> TYLENOL[®] Reliever ARTHRITIS PAIN</p> <p><i>Contains</i> Acetaminophen Extended Release FOR THE TEMPORARY RELIEF OF MINOR ARTHRITIS PAIN</p> <p>100 Caplets* - 650 mg EACH *Capsule-Shaped Tablets</p>	<p>Directions ■ do not take more than directed (see overdose warning)</p> <p>Adults: ■ take 2 caplets every 8 hours with water ■ swallow whole - do not crush, chew or dissolve ■ do not take more than 6 caplets in 24 hours ■ do not use for more than 10 days unless directed by a doctor</p> <p>Under 18 years of age: ■ ask a doctor</p> <p>Other information ■ do not use if neck wrap or foil inner seal imprinted with "Safety Seal[®]" is broken ■ store at 20 - 25°C (68 - 77°F) ■ avoid excessive heat 40°C (104°F)</p> <p>Inactive ingredients corn starch, hydroxyethyl cellulose, hypromellose, magnesium stearate, microcrystalline cellulose, povidone, powdered cellulose, pregelatinized starch, sodium starch glycolate, titanium dioxide, triacetin</p> <p>Questions or comments? call toll-free 1-877-TYLENOL (1-877-695-3645)</p>

NDC 50580-112-10
 TYLENOL[®] Reliever
 ARTHRITIS PAIN
 100 Caplets* - 650 mg EACH
 *Capsule-Shaped Tablets
 © 2002
 PAIN RELIEVER
 U.S. PAT. NO. 4,808,822
 www.tylenol.com
 TYLENOL[®] is a registered trademark of Parke-Davis Inc.
 Parke-Davis Inc., a subsidiary of Pharmacia Corporation
 2455 Ridge Pike, North Wales, PA 19381 USA

Labeling Format Information:			
Fonts: Helvetica regular, bold and bold italic.			
Drug Facts:	NA	Leading:	5 pt
Header:	6 pt	Bullets:	4 pt
Subheader:	6 pt	Barlines:	NA
Body Text:	5 pt	Hairlines:	NA
Drug Facts (continued):	NA		
Horizontal Scale: 80%		Average Kerning: 0	

LBL TY ARTH ER CPLT 100 3004009 B-2
OCTOBER 16, 2002