


EV CODE AREA TO BE
COATING AND COPY FREE

724



Drug Facts

Read the directions and warnings before taking this product. KEEP THE PRODUCT INFORMATION SHEET. IT CONTAINS IMPORTANT INFORMATION.

Active ingredient (in each tablet) Cimetidine 200 mg.....

Purpose Acid reducer

Uses • relieves heartburn associated with acid indigestion and sour stomach
• prevents heartburn associated with acid indigestion and sour stomach brought on by eating or drinking certain food and beverages

Warnings

Allergy alert: Do not use if you are allergic to cimetidine or other acid reducers

Do not use

• if you have trouble or pain swallowing food, vomiting with blood, or bloody or black stools. These may be signs of a serious condition. See your doctor.
• with other acid reducers

Ask a doctor before use if you have

• frequent chest pain • frequent wheezing, particularly with heartburn
• unexplained weight loss • nausea or vomiting
• stomach pain
• had heartburn over 3 months. This may be a sign of a more serious condition.
• heartburn with lightheadedness, sweating or dizziness
• chest pain or shoulder pain with shortness of breath; sweating; pain spreading to arms, neck or shoulders; or lightheadedness

Ask a doctor or pharmacist before use if you are taking

• theophylline (oral asthma medicine) • warfarin (blood thinning medicine)
• phenytoin (seizure medicine)

If you are not sure you are taking one of these medicines, talk to your doctor or pharmacist.

Stop use and ask a doctor if

• your heartburn continues or worsens • stomach pain continues
• you need to take this product for more than 14 days

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

• adults and children 12 years and over:
• to relieve symptoms, swallow 1 tablet with a glass of water
• to prevent symptoms, swallow 1 tablet with a glass of water right before or any time up to 30 minutes before eating food or drinking beverages that cause heartburn
• do not take more than 2 tablets in 24 hours
• children under 12 years: ask a doctor

Other information • store at 15-30°C (59-86°F)

Inactive ingredients cellulose, corn starch, hypromellose, magnesium stearate, polyethylene glycol, polysorbate 80, povidone, sodium lauryl sulfate, sodium starch glycolate, titanium dioxide

Questions or comments?

call toll-free 1-800-482-4394 weekdays



3 0766-5016-06 4

KNOCKOUT FOR GLUE

"LOT" & "EXP" TO BE PREPRINTED
CODE IS DEBOSSSED

Each tablet individually sealed in a blister pack. Do not use if blister or foil is open or torn.

LOT EXP.

9046X

SAVE UP TO \$4.00
(money-saving offer inside)

Just **ONE TABLET**
RELIEVES and **PREVENTS**
Heartburn and Acid Indigestion

Tagamet
Cimetidine Tablets 200 mg/Acid Reducer **HB 200**



6 TABLETS (6 DOSES)

Take **ANY TIME** you need it:

- Before Meal
- During Meal
- After Meal

Tagamet
HB 200

SAVE UP TO \$4.00
(money-saving offer inside)


Just **ONE TABLET**
RELIEVES and **PREVENTS**
Heartburn and Acid Indigestion

Tagamet
HB 200

Just **ONE TABLET**
RELIEVES and **PREVENTS**
Heartburn and Acid Indigestion

Distributed by:
GlaxoSmithKline Consumer Healthcare, L.P.
Moon Twp., PA 15108, Made in the U.S.A.
©2004 GlaxoSmithKline

724



EV CODE AREA TO BE
COATING AND COPY FREE