



Rx only

AZACTAM®

(aztreonam injection)



in Galaxy® Plastic Container (PL 2040) for Intravenous Use

DESCRIPTION

AZACTAM® (aztreonam injection) contains the active ingredient, aztreonam, a monobactam. It was originally isolated from *Chromobacterium violaceum*. It is a synthetic bactericidal antibiotic.

The monobactams, having a unique monocyclic beta-lactam nucleus, are structurally different from other beta-lactam antibiotics (e.g., penicillins, cephalosporins, cephamycins). The sulfonic acid substituent in the 1-position of the ring activates the beta-lactam moiety; an aminothiazolyl oxime side chain in the 3-position and a methyl group in the 4-position confer the specific antibacterial spectrum and beta-lactamase stability.

Aztreonam is designated chemically as (Z)-2-[[[(2-amino-4-thiazolyl)[[(2S,-3S)-2-methyl-4-oxo-1-sulfo-3-azetidiny]carbamoyl]methylene]amino]oxy]-2-methylpropionic acid. Structural formula:



AZACTAM (aztreonam injection) in the Galaxy plastic container (PL 2040) is a frozen, iso-osmotic, sterile, sodium-free, nonpyrogenic intravenous solution. Each 50 mL of solution contains 1 g, or 2 g aztreonam with approximately 1.7 g, or 700 mg dextrose hydrous, USP added to adjust osmolality, and approximately 780 mg, or 1.6 g of arginine added for pH adjustment, respectively. Thawed solutions have a pH in the range of 4.5 to 7.5. The solution is for intravenous administration following thawing at room temperature or under refrigeration.


This Galaxy container is fabricated from a specially designed multilayer plastic (PL 2040). Solutions are in contact with the polyethylene layer of this container and can leach out certain chemical components of the plastic in very small amounts within the expiration period. The suitability of the plastic has been confirmed in tests in animals according to the USP biological tests for plastic containers as well as by tissue culture toxicity studies.

CLINICAL PHARMACOLOGY

Single 30-minute intravenous infusions of 500 mg, 1 g and 2 g doses of AZACTAM in healthy subjects produced peak serum levels of 54, 90 and 204 $\mu\text{g/mL}$, respectively, immediately after administration; at eight hours, serum levels were 1, 3 and 6 $\mu\text{g/mL}$, respectively (Figure 1). Single 3-minute intravenous injections of the same doses resulted in serum levels of 58, 125 and 242 $\mu\text{g/mL}$ at five minutes following completion of injection.

Serum concentrations of aztreonam following completion of single intravenous infusions of 500 mg, 1 g, and 2 g doses are depicted in Figure 1.

FIGURE 1



The serum levels of aztreonam following single 500 mg, 1 g or 2 g intravenous doses of AZACTAM (aztreonam injection) exceed the MIC_{90} for *Neisseria* sp., *H. influenzae* and most genera of the *Enterobacteriaceae* for eight hours (for *Enterobacter* sp., the eight hour serum levels exceed the MIC for 80 percent of strains). For *P. aeruginosa*, a single 2 g intravenous dose produces serum levels that exceed the MIC_{90} for approximately four to six hours. All of the above doses of AZACTAM result in average urine levels of aztreonam that exceed the MIC_{90} for the same pathogens for up to 12 hours.

When aztreonam pharmacokinetics were assessed for adult and pediatric patients, they were found to be comparable (down to 9 months old). The serum half-life of aztreonam averaged 1.7 hours (1.5 to 2.0) in subjects with normal renal function, independent of the

dose. In healthy subjects, based on a 70 kg person, the serum clearance was 91 mL/min and renal clearance was 56 mL/min; the apparent mean volume of distribution at steady-state averaged 12.6 liters, approximately equivalent to extracellular fluid volume.

In a study of healthy elderly male subjects (65 to 75 years of age), the average elimination half-life of aztreonam was slightly longer than in young healthy males.

In patients with impaired renal function, the serum half-life of aztreonam is prolonged (see **DOSAGE AND ADMINISTRATION, Renal Impairment in Adult Patients**). The serum half-life of aztreonam is only slightly prolonged in patients with hepatic impairment since the liver is a minor pathway of excretion.

Average urine concentrations of aztreonam were approximately 1100, 3500 and 6600 µg/mL within the first two hours following single 500 mg, 1 g and 2 g intravenous doses of AZACTAM (30-minute infusions), respectively. The range of average concentrations for aztreonam in the 8 to 12 hour urine specimens in these studies was 25 to 120 µg/mL. In healthy subjects, aztreonam is excreted in the urine about equally by active tubular secretion and glomerular filtration. Approximately 60 to 70 percent of an intravenous dose was recovered in the urine by eight hours. Urinary excretion of a single intravenous dose was essentially complete by 12 hours after injection. About 12 percent of a single intravenous radiolabeled dose was recovered in the feces. Unchanged aztreonam and the inactive beta-lactam ring hydrolysis product of aztreonam were present in feces and urine.

Intravenous administration of a single 500 mg or 1 g dose of AZACTAM every eight hours for seven days to healthy subjects produced no apparent accumulation of aztreonam or modification of its disposition characteristics; serum protein binding averaged 56 percent and was independent of dose.

Renal function was monitored in healthy subjects given aztreonam; standard tests (serum creatinine, creatinine clearance, BUN, urinalysis and total urinary protein excretion) as well as special tests (excretion of N-acetyl-β-glucosaminidase, alanine aminopeptidase and β₂-microglobulin) were used. No abnormal results were obtained.

Aztreonam achieves measurable concentrations in the following body fluids and tissues:

EXTRAVASCULAR CONCENTRATIONS OF AZTREONAM					
AFTER A SINGLE INTRAVENOUS DOSE¹					
Fluid or Tissue	Dose (g)	Route	Hours Postin- jection	Number of Patients	Mean Concentration (µg/mL or µg/g)
Fluids					
bile	1	IV	2	10	39
blister fluid	1	IV	1	6	20
bronchial secretion	2	IV	4	7	5
cerebrospinal fluid (inflamed meninges)	2	IV	0.9-4.3	16	3
pericardial fluid	2	IV	1	6	33
pleural fluid	2	IV	1.1-3.0	3	51
synovial fluid	2	IV	0.8-1.9	11	83
Tissues					
atrial appendage	2	IV	0.9-1.6	12	22
endometrium	2	IV	0.7-1.9	4	9
fallopian tube	2	IV	0.7-1.9	8	12
fat	2	IV	1.3-2.0	10	5
femur	2	IV	1.0-2.1	15	16
gallbladder	2	IV	0.8-1.3	4	23
kidney	2	IV	2.4-5.6	5	67
large intestine	2	IV	0.8-1.9	9	12
liver	2	IV	0.9-2.0	6	47
lung	2	IV	1.2-2.1	6	22
myometrium	2	IV	0.7-1.9	9	11
ovary	2	IV	0.7-1.9	7	13
skeletal muscle	2	IV	0.3-0.7	6	16
skin	2	IV	0.0-1.0	8	25
sternum	2	IV	1	6	6

¹Tissue penetration is regarded as essential to therapeutic efficacy, but specific tissue levels have not been correlated with specific therapeutic effects.

The concentration of aztreonam in saliva at 30 minutes after a single 1 g intravenous dose (9 patients) was 0.2 µg/mL; in breast milk at two hours after a single 1 g intravenous dose (6 patients), 0.2 µg/mL; in amniotic fluid at six to eight hours after a single 1 g intravenous dose (5 patients), 2 µg/mL. The concentration of aztreonam in peritoneal fluid obtained one to six hours after multiple 2 g intravenous doses ranged between 12 and 90 µg/mL in 7 of 8 patients studied.

Aztreonam given intravenously rapidly reaches therapeutic concentrations in peritoneal dialysis fluid; conversely, aztreonam given intraperitoneally in dialysis fluid rapidly produces therapeutic serum levels.

Concomitant administration of probenecid or furosemide and aztreonam causes clinically insignificant increases in the serum levels of aztreonam. Single-dose intravenous pharmacokinetic studies have not shown any significant interaction between aztreonam and concomitantly administered gentamicin, nafcillin sodium, cephadrine, clindamycin or metronidazole. No reports of disulfiram-like reactions with alcohol ingestion have been noted; this is not unexpected since aztreonam does not contain a methyl-tetrazole side chain.

Microbiology

Aztreonam exhibits potent and specific activity *in vitro* against a wide spectrum of gram-negative aerobic pathogens including *Pseudomonas aeruginosa*. The bactericidal action of aztreonam results from the inhibition of bacterial cell wall synthesis due to a high affinity of aztreonam for penicillin binding protein 3 (PBP3). Aztreonam, unlike the majority of beta-lactam antibiotics, does not induce beta-lactamase activity and its molecular structure confers a high degree of resistance to hydrolysis by beta-lactamases (i.e., penicillinases and cephalosporinases) produced by most gram-negative and gram-positive pathogens; it is, therefore, usually active against gram-negative aerobic microorganisms that are resistant to antibiotics hydrolyzed by beta-lactamases. It is active against many strains that are multiply-resistant to other antibiotics, such as certain cephalosporins, penicillin, and aminoglycosides. Aztreonam maintains its antimicrobial activity over a pH range of 6 to 8 *in vitro*, as well as in the presence of human serum and under anaerobic conditions.

Aztreonam has been shown to be active against most strains of the following microorganisms, both *in vitro* and in clinical infections as described in the **INDICATIONS AND USAGE** section.

Aerobic gram-negative microorganisms:

Citrobacter species, including *C. freundii*

Enterobacter species, including *E. cloacae*

Escherichia coli

Haemophilus influenzae (including ampicillin-resistant and other penicillinase-producing strains)

Klebsiella oxytoca

Klebsiella pneumoniae

Proteus mirabilis

Pseudomonas aeruginosa
Serratia species, including *S. marcescens*

The following *in vitro* data are available, **but their clinical significance is unknown.**

Aztreonam exhibits *in vitro* minimal inhibitory concentrations (MIC's) of 8 µg/mL or less against most (≥90%) strains of the following microorganisms; however, the safety and effectiveness of aztreonam in treating clinical infections due to these microorganisms have not been established in adequate and well-controlled clinical trials.

Aerobic gram-negative microorganisms:

Aeromonas hydrophila
Morganella morganii
Neisseria gonorrhoeae (including penicillinase-producing strains)
Pasteurella multocida
Proteus vulgaris
Providencia stuartii
Providencia rettgeri
Yersinia enterocolitica

Aztreonam and aminoglycosides have been shown to be synergistic *in vitro* against most strains of *P. aeruginosa*, many strains of *Enterobacteriaceae*, and other gram-negative aerobic bacilli.

Alterations of the anaerobic intestinal flora by broad spectrum antibiotics may decrease colonization resistance, thus permitting overgrowth of potential pathogens, e.g., *Candida* and *Clostridium* species. Aztreonam has little effect on the anaerobic intestinal microflora in *in vitro* studies. *Clostridium difficile* and its cytotoxin were not found in animal models following administration of aztreonam. (See **ADVERSE REACTIONS**, *Gastrointestinal*.)

Susceptibility Tests

Dilution Techniques: Quantitative methods are used to determine antimicrobial minimal inhibitory concentrations (MIC's). These MIC's provide estimates of the susceptibility of bacteria to antimicrobial compounds. The MIC's should be determined using a standardized procedure. Standardized procedures are based on a dilution method¹ (broth or agar) or equivalent with standardized inoculum concentrations and standardized concentrations of aztreonam powder. The MIC values should be interpreted according to the following criteria:

For testing aerobic microorganisms other than *Haemophilus influenzae*:

<u>MIC (µg/mL)</u>	<u>Interpretation</u>
≤8	Susceptible (S)
16	Intermediate (I)
≥32	Resistant (R)

When testing *Haemophilus influenzae*^a:

<u>MIC (µg/mL)</u>	<u>Interpretation^b</u>
≤2	Susceptible (S)

a. Interpretative criteria applicable only to tests performed by broth microdilution method using *Haemophilus* Test Medium (HTM)¹.

b. The current absence of data on resistant strains precludes defining any categories other than “Susceptible”. Strains yielding MIC results suggestive of a “nonsusceptible” category should be submitted to a reference laboratory for further testing.

A report of “Susceptible” indicates that the pathogen is likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable. A report of “Intermediate” indicates that the result should be considered equivocal, and, if the microorganism is not fully susceptible to alternative, clinically feasible drugs, the test should be repeated. This category implies possible clinical applicability in body sites where the drug is physiologically concentrated or in situations where high dosage of drug can be used. This category also provides a buffer zone which prevents small uncontrolled technical factors from causing major discrepancies in interpretation. A report of “Resistant” indicates that the pathogen is not likely to be inhibited if the antimicrobial compound in the blood reaches the concentrations usually achievable; other therapy should be selected.

Standardized susceptibility test procedures require the use of laboratory control microorganisms to control the technical aspects of the laboratory procedures. Standard aztreonam powder should provide the following MIC values:

<u>Microorganism</u>	<u>MIC (µg/mL)</u>
<i>Escherichia coli</i> ATCC 25922	0.06-0.25
<i>Haemophilus influenzae</i> ^a ATCC 49247	0.12-0.5
<i>Pseudomonas aeruginosa</i> ATCC 27853	2.0-8.0

a. Range applicable only to tests performed by broth microdilution method using *Haemophilus* Test Medium (HTM)¹.

Diffusion Techniques: Quantitative methods that require measurement of zone diameters also provide reproducible estimates of the susceptibility of bacteria to antimicrobial compounds. One such standardized procedure² requires the use of standardized inoculum concentrations. This procedure uses paper disks impregnated with 30-µg aztreonam to test the susceptibility of microorganisms to aztreonam.

Reports from the laboratory providing results of the standard single-disk susceptibility test with a 30-µg aztreonam disk should be interpreted according to the following criteria:

For testing aerobic microorganisms other than *Haemophilus influenzae*:

<u>Zone diameter (mm)</u>	<u>Interpretation</u>
≥22	Susceptible (S)
16 - 21	Intermediate (I)
≤15	Resistant (R)

When testing *Haemophilus influenzae*^a:

<u>Zone diameter (mm)</u>	<u>Interpretation^b</u>
≥26	Susceptible (S)

a. Interpretative criteria applicable only to tests performed by disk diffusion method using *Haemophilus* Test Medium (HTM)².

b. The current absence of data on resistant strains precludes defining any categories other than "Susceptible". Strains yielding zone diameter results suggestive of a "nonsusceptible" category should be submitted to a reference laboratory for further testing.

Interpretation should be as stated above for results using dilution techniques. Interpretation involves correlation of the diameter obtained in the disk test with the MIC for aztreonam.

As with standardized dilution techniques, diffusion methods require the use of laboratory control microorganisms that are used to control the technical aspects of the laboratory procedures. For the diffusion technique, the 30-μg aztreonam disk should provide the following zone diameters in these laboratory test quality control strains.

<u>Microorganism</u>	<u>Zone diameter (mm)</u>
<i>Escherichia coli</i> ATCC 25922	28-36 mm
<i>Haemophilus influenzae</i> ^a ATCC 49247	30-38 mm
<i>Pseudomonas aeruginosa</i> ATCC 27853	23-29 mm

a. Range applicable only to tests performed by dish diffusion method using *Haemophilus* Test Medium (HTM)².

INDICATIONS AND USAGE

Before initiating treatment with AZACTAM, appropriate specimens should be obtained for isolation of the causative organism(s) and for determination of susceptibility to aztreonam. Treatment with AZACTAM may be started empirically before results of the susceptibility testing are available; subsequently, appropriate antibiotic therapy should be continued.

AZACTAM (aztreonam injection) is indicated for the treatment of the following infections caused by susceptible gram-negative microorganisms:

Urinary Tract Infections (complicated and uncomplicated), including pyelonephritis and cystitis (initial and recurrent) caused by *Escherichia coli*, *Klebsiella pneumoniae*, *Proteus mirabilis*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Klebsiella oxytoca**, *Citrobacter species** and *Serratia marcescens**.

Lower Respiratory Tract Infections, including pneumonia and bronchitis caused by *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Haemophilus*

influenzae, *Proteus mirabilis*, *Enterobacter* species and *Serratia marcescens**.

Septicemia caused by *Escherichia coli*, *Klebsiella pneumoniae*, *Pseudomonas aeruginosa*, *Proteus mirabilis**, *Serratia marcescens** and *Enterobacter* species.

Skin and Skin-Structure Infections, including those associated with postoperative wounds, ulcers and burns caused by *Escherichia coli*, *Proteus mirabilis*, *Serratia marcescens*, *Enterobacter* species, *Pseudomonas aeruginosa*, *Klebsiella pneumoniae* and *Citrobacter* species*.

Intra-abdominal Infections, including peritonitis caused by *Escherichia coli*, *Klebsiella* species including *K. pneumoniae*, *Enterobacter* species including *E. cloacae**, *Pseudomonas aeruginosa*, *Citrobacter* species* including *C. freundii** and *Serratia* species* including *S. marcescens**.

Gynecologic Infections, including endometritis and pelvic cellulitis caused by *Escherichia coli*, *Klebsiella pneumoniae**, *Enterobacter* species* including *E. cloacae** and *Proteus mirabilis**.

AZACTAM (aztreonam injection) is indicated for adjunctive therapy to surgery in the management of infections caused by susceptible organisms, including abscesses, infections complicating hollow viscus perforations, cutaneous infections and infections of serous surfaces. AZACTAM is effective against most of the commonly encountered gram-negative aerobic pathogens seen in general surgery.

*Efficacy for this organism in this organ system was studied in fewer than ten infections.

Concurrent Therapy

Concurrent initial therapy with other antimicrobial agents and AZACTAM is recommended before the causative organism(s) is known in seriously ill patients who are also at risk of having an infection due to gram-positive aerobic pathogens. If anaerobic organisms are also suspected as etiologic agents, therapy should be initiated using an anti-anaerobic agent concurrently with AZACTAM (see **DOSAGE AND ADMINISTRATION**). Certain antibiotics (e.g., cefoxitin, imipenem) may induce high levels of beta-lactamase *in vitro* in some gram-negative aerobes such as *Enterobacter* and *Pseudomonas* species, resulting in antagonism to many beta-lactam antibiotics including aztreonam. These *in vitro* findings suggest that such beta-lactamase inducing antibiotics not be used concurrently with aztreonam. Following identification and susceptibility testing of the causative organism(s), appropriate antibiotic therapy should be continued.

CONTRAINDICATIONS

This preparation is contraindicated in patients with known hypersensitivity to aztreonam or any other component in the formulation.

WARNINGS

Both animal and human data suggest that AZACTAM is rarely cross-reactive with other beta-lactam antibiotics and weakly immunogenic. Treatment with aztreonam can result in hypersensitivity reactions in patients with or without prior exposure. (See **CONTRAINDICATIONS**.)

Careful inquiry should be made to determine whether the patient has any history of hypersensitivity reactions to any allergens.

While cross-reactivity of aztreonam with other beta-lactam antibiotics is rare, this drug should be administered with caution to any patient with a history of hypersensitivity to beta-lactams (e.g., penicillins, cephalosporins, and/or carbapenems). Treatment with aztreonam can result in hypersensitivity reactions in patients with or without prior exposure to aztreonam. If an allergic reaction to aztreonam occurs, discontinue the drug and institute supportive treatment as appropriate (e.g., maintenance of ventilation, pressor amines, antihistamines, corticosteroids). Serious hypersensitivity reactions may require epinephrine and other emergency measures. (See **ADVERSE REACTIONS**.)

Pseudomembranous colitis has been reported with nearly all antibacterial agents, including aztreonam, and may range in severity from mild to life-threatening. Therefore, it is important to consider this diagnosis in patients who present with diarrhea subsequent to the administration of antibacterial agents.

Treatment with antibacterial agents alters the normal flora of the colon and may permit overgrowth of *Clostridia*. Studies indicate that a toxin produced by *Clostridium difficile* is one primary cause of "antibiotic-associated colitis."

After the diagnosis of pseudomembranous colitis has been established, therapeutic measures should be initiated. Mild cases of pseudomembranous colitis usually respond to drug discontinuation alone. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial drug clinically effective against *C. difficile* colitis.

Rare cases of toxic epidermal necrolysis have been reported in association with aztreonam in patients undergoing bone marrow transplant with multiple risk factors including sepsis, radiation therapy and other concomitantly administered drugs associated with toxic epidermal necrolysis.

PRECAUTIONS

General

In patients with impaired hepatic or renal function, appropriate monitoring is recommended during therapy.

If an aminoglycoside is used concurrently with aztreonam, especially if high dosages of the former are used or if therapy is prolonged, renal function should be monitored because of the potential nephrotoxicity and ototoxicity of aminoglycoside antibiotics.

The use of antibiotics may promote the overgrowth of nonsusceptible organisms, including gram-positive organisms (*Staphylococcus aureus* and *Streptococcus faecalis*) and fungi. Should superinfection occur during therapy, appropriate measures should be taken.

Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenicity studies in animals have not been performed.

Genetic toxicology studies performed *in vivo* and *in vitro* with aztreonam in several standard laboratory models revealed no evidence of mutagenic potential at the chromosomal or gene level.

Two-generation reproduction studies in rats at daily doses up to 20 times the maximum recommended human dose, prior to and during gestation and lactation, revealed no evidence of impaired fertility. There was a slightly reduced survival rate during the lactation period in the offspring of rats that received the highest dosage, but not in offspring of rats that received five times the maximum recommended human dose.

Pregnancy Category B

Aztreonam crosses the placenta and enters the fetal circulation.

Studies in pregnant rats and rabbits, with daily doses up to 15 and 5 times, respectively, the maximum recommended human dose, revealed no evidence of embryo- or fetotoxicity or teratogenicity. No drug induced changes were seen in any of the maternal, fetal, or neonatal parameters that were monitored in rats receiving 15 times the maximum recommended human dose of aztreonam during late gestation and lactation.

There are no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, aztreonam should be used during pregnancy only if clearly needed.

Nursing Mothers

Aztreonam is excreted in breast milk in concentrations that are less than 1 percent of concentrations determined in simultaneously obtained maternal serum; consideration should be given to temporary discontinuation of nursing and use of formula feedings.

Pediatric Use

The safety and effectiveness of intravenous AZACTAM have been established in the age groups 9 months to 16 years. Use of AZACTAM in these age groups is supported by evidence from adequate and well-controlled studies of AZACTAM in adults with additional efficacy, safety, and pharmacokinetic data from noncomparative clinical studies in pediatric patients. Sufficient data are not available for pediatric patients under 9 months of age or for the following treatment indications/pathogens: septicemia and skin and skin-structure infections (where the skin infection is believed or known to be due to *H. influenzae* type b). In pediatric patients with cystic fibrosis, higher doses of AZACTAM may be warranted. (See **CLINICAL PHARMACOLOGY, DOSAGE AND ADMINISTRATION**, and **CLINICAL STUDIES**.)

ADVERSE REACTIONS

Local reactions (e.g., phlebitis/thrombophlebitis; discomfort/swelling) following IV administration occurred at rates of approximately 1.9 percent.

Systemic reactions (considered to be related to therapy or of uncertain etiology) occurring at an incidence of 1 to 1.3 percent include diarrhea, nausea and/or vomiting, and rash. Reactions occurring at an incidence of less than 1 percent are listed within each body system in order of decreasing severity:

Hypersensitivity—anaphylaxis, angioedema, bronchospasm.

Hematologic—pancytopenia, neutropenia, thrombocytopenia, anemia, eosinophilia, leukocytosis, thrombocytosis.

Gastrointestinal—abdominal cramps; rare cases of *C. difficile*-associated diarrhea, including pseudomembranous colitis, or gastrointestinal bleeding have been reported. Onset of pseudomembranous colitis symptoms may occur during or after antibiotic treatment. (See **WARNINGS**.)

Dermatologic—toxic epidermal necrolysis (see **WARNINGS**), purpura, erythema multiforme, exfoliative dermatitis, urticaria, petechiae, pruritus, diaphoresis.

Cardiovascular—hypotension, transient ECG changes (ventricular bigeminy and PVC), flushing.

Respiratory—wheezing, dyspnea, chest pain.

Hepatobiliary—hepatitis, jaundice.

Nervous System—seizure, confusion, vertigo, paresthesia, insomnia, dizziness.

Musculoskeletal—muscular aches.

Special Senses—tinnitus, diplopia, mouth ulcer, altered taste, numb tongue, sneezing, nasal congestion, halitosis.

Other—vaginal candidiasis, vaginitis, breast tenderness.

Body as a Whole—weakness, headache, fever, malaise.

Pediatric Adverse Reactions

Of the 612 pediatric patients who were treated with AZACTAM in clinical trials, less than 1% required discontinuation of therapy due to adverse events. The following systemic adverse events, regardless of drug relationship, occurred in at least 1% of treated patients in domestic clinical trials: rash (4.3%), diarrhea (1.4%), and fever (1.0%). These adverse events were comparable to those observed in adult clinical trials.

In 343 pediatric patients receiving intravenous therapy, the following local reactions were noted: pain (12%), erythema (2.9%), induration (0.9%), and phlebitis (2.1%). In the US patient population, pain occurred in 1.5% of patients, while each of the remaining three local reactions had an incidence of 0.5%.

The following laboratory adverse events, regardless of drug relationship, occurred in at least 1% of treated patients: increased eosinophils (6.3%), increased platelets (3.6%), neutropenia (3.2%), increased AST (3.8%), increased ALT (6.5%), and increased serum creatinine (5.8%).

In US pediatric clinical trials, neutropenia (absolute neutrophil count less than 1000/mm³) occurred in 11.3% of patients (8/71) younger than 2 years receiving 30 mg/kg q6h. AST and ALT elevations to greater than 3 times the upper limit of normal were noted in 15-20% of patients aged 2 years or above receiving 50 mg/kg q6h. The increased frequency of these reported laboratory adverse events may be due to either increased severity of illness treated or higher doses of AZACTAM administered.

Adverse Laboratory Changes

Adverse laboratory changes without regard to drug relationship that were reported during clinical trials were:

Hepatic—elevations of AST (SGOT), ALT (SGPT), and alkaline phosphatase; signs or

symptoms of hepatobiliary dysfunction occurred in less than 1 percent of recipients (see above).

Hematologic—increases in prothrombin and partial thromboplastin times, positive Coombs' test.

Renal—increases in serum creatinine.

OVERDOSAGE

If necessary, aztreonam may be cleared from the serum by hemodialysis and/or peritoneal dialysis.

DOSAGE AND ADMINISTRATION

Dosage in Adult Patients

AZACTAM (aztreonam injection), an intravenous solution in Galaxy plastic containers (PL 2040), is intended for intravenous use only. Dosage should be determined by susceptibility of the causative organisms, severity and site of infection, and the condition of the patient.

The intravenous route is recommended for patients with bacterial septicemia, localized parenchymal abscess (e.g., intra-abdominal abscess), peritonitis or other severe systemic or life-threatening infections.

The duration of therapy depends on the severity of infection. Generally, AZACTAM should be continued for at least 48 hours after the patient becomes asymptomatic or evidence of bacterial eradication has been obtained. Persistent infections may require treatment for several weeks. Doses smaller than those indicated should not be used.

Renal Impairment in Adult Patients

Prolonged serum levels of aztreonam may occur in patients with transient or persistent renal insufficiency. Therefore, the dosage of AZACTAM should be halved in patients with estimated creatinine clearances between 10 and 30 mL/min/1.73 m² after an initial loading dose of 1 g or 2 g.

When only the serum creatinine concentration is available, the following formula (based on sex, weight, and age of the patient) may be used to approximate the creatinine clearance (Clcr). The serum creatinine should represent a steady state of renal function.

$$\text{Males: Clcr} = \frac{\text{weight (kg)} \times (140 - \text{age})}{72 \times \text{serum creatinine (mg/dL)}}$$

Females: 0.85 x above value

In patients with severe renal failure (creatinine clearance less than 10 mL/min/1.73 m²), such as those supported by hemodialysis, the usual dose of 500 mg, 1 g or 2 g should be given initially. The maintenance dose should be one-fourth of the usual initial dose given at the usual fixed interval of 6, 8 or 12 hours. For serious or life-threatening infections, in addition to the maintenance doses, one-eighth of the initial dose should be given after each hemodialysis session.

Dosage In The Elderly

Renal status is a major determinant of dosage in the elderly; these patients in particular may have diminished renal function. Serum creatinine may not be an accurate determinant of renal status. Therefore, as with all antibiotics eliminated by the kidneys, estimates of creatinine clearance should be obtained, and appropriate dosage modifications made if necessary.

Dosage in Pediatric Patients

AZACTAM should be administered intravenously to pediatric patients with normal renal function. There are insufficient data regarding intramuscular administration to pediatric patients or dosing in pediatric patients with renal impairment. (See **PRECAUTIONS: Pediatric Use.**)

AZACTAM DOSAGE GUIDELINES

Type of Infection	Dose	Frequency (hours)
ADULTS*		
Urinary tract infections	500 mg or 1 g	8 or 12
Moderately severe systemic infections	1 g or 2 g	8 or 12
Severe systemic or life-threatening infections	2 g	6 or 8
*Maximum recommended dose is 8 g per day		
PEDIATRIC PATIENTS**		
Mild to moderate infections	30 mg/kg	8
Moderate to severe infections	30 mg/kg	6 or 8
**Maximum recommended dose is 120 mg/kg/day		

Because of the serious nature of infections due to *Pseudomonas aeruginosa*, dosage of 2 g every six or eight hours is recommended, at least upon initiation of therapy, in systemic infections caused by this organism in adults.

CLINICAL STUDIES

A total of 612 pediatric patients aged 1 month to 12 years were enrolled in uncontrolled clinical trials of aztreonam in the treatment of serious gram-negative infections, including urinary tract, lower respiratory tract, skin and skin-structure, and intra-abdominal infections.

Directions for Use of AZACTAM (aztreonam injection) in Galaxy Plastic Container (PL 2040).

AZACTAM (aztreonam injection) in Galaxy plastic container (PL 2040) is to be administered as an intermittent intravenous infusion only.

Storage

Store in a freezer capable of maintaining a temperature of -20° C (-4° F).

Thawing of Plastic Containers

Thaw frozen container at room temperature, 25° C (77° F) or in a refrigerator, 2° to 8° C (36° to 46° F). After thawing is complete, invert the container to assure a well-mixed solution. **(DO NOT FORCE THAW BY IMMERSION IN WATER BATHS OR BY MICROWAVE IRRADIATION.)**

Check for minute leaks by squeezing container firmly. If leaks are detected, discard solution as sterility may be impaired.

The container should be visually inspected. Thawed solutions should not be used unless clear; solutions will be colorless to yellow. Components of the solution may precipitate in the frozen state and will dissolve upon reaching room temperature with little or no agitation. If after visual inspection the solution remains discolored, cloudy, or if an insoluble precipitate is noted or if any seals or outlet ports are not intact, the container should be discarded.

DO NOT ADD SUPPLEMENTARY MEDICATION.

The thawed solution in Galaxy plastic container (PL 2040) remains chemically stable for either 14 days under refrigeration (2° to 8° C/36° to 46° F) or for 48 hours at room temperature (25° C/77° F). **DO NOT REFREEZE THAWED ANTIBIOTICS.**

Preparation for Intravenous Administration (Use aseptic technique)

1. Suspend container(s) from eyelet support.
2. Remove protector from outlet port at bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

Additives or other medication should not be added to AZACTAM (aztreonam injection) in Galaxy plastic container (PL 2040) or infused simultaneously through the same intravenous line. If the same intravenous line is used for sequential infusion of several different drugs, it should be flushed before and after infusion of AZACTAM (aztreonam injection) in Galaxy plastic container (PL 2040) with an infusion solution compatible with AZACTAM (aztreonam injection) in Galaxy plastic container (PL 2040)* and any other drug(s) administered via this common line.

It is recommended that the intravenous administration apparatus be replaced at least once every 48 hours.

CAUTION: Do not use plastic containers in series connections. Such use could result in an embolism due to residual air being drawn from the primary container before administration of the fluid from the secondary container is complete.

Intravenous Administration

Infusion of AZACTAM (aztreonam injection) in Galaxy plastic container (PL 2040) should be completed within a 20 to 60 minute period. The plastic container is a single-dose unit; discard any unused portion remaining in the container.

*The following infusion solutions are compatible with AZACTAM (aztreonam injection) in Galaxy plastic container (PL 2040):

Sodium Chloride Injection USP, 0.9%
Ringer's Injection USP
Lactated Ringer's Injection USP
Dextrose Injection USP, 5% or 10%
Dextrose and Sodium Chloride Injection USP, 5%:0.9%,
5%:0.45% or 5%:0.2%
Sodium Lactate Injection USP (M/6 Sodium Lactate)
Ionosol® B and 5% Dextrose
Isolyte® E
Isolyte® E with 5% Dextrose
Isolyte® M with 5% Dextrose
Normosol®-R
Normosol®-R and 5% Dextrose
Normosol®-M and 5% Dextrose
Mannitol Injection USP, 5% or 10%
Lactated Ringer's and 5% Dextrose Injection
Plasma-Lyte® M and 5% Dextrose
10% Travert® Injection
10% Travert® and Electrolyte No. 1 Injection
10% Travert® and Electrolyte No. 2 Injection
10% Travert® and Electrolyte No. 3 Injection

HOW SUPPLIED

AZACTAM® (aztreonam injection) in Galaxy® plastic container (PL 2040) is supplied as a frozen, 50 mL single-dose intravenous solution as follows:

1-g aztreonam/50 mL container:

Packages of 24 (NDC 0003-2231-01)

2-g aztreonam/50 mL container:

Packages of 24 (NDC 0003-2241-01)

Store at or below -20° C (-4° F) [See **Directions for Use of AZACTAM® (aztreonam injection) in Galaxy® Plastic Container (PL 2040)**].

