

**DIAMOX® SEQUELS® (Acetazolamide Extended-Release Capsules)**

**Rx only**

**DESCRIPTION**

DIAMOX SEQUELS (Acetazolamide Extended-Release Capsules) are an inhibitor of the enzyme carbonic anhydrase.

DIAMOX is a white to faintly yellowish white crystalline, odorless powder, weakly acidic, very slightly soluble in water and slightly soluble in alcohol. The chemical name for DIAMOX is N-(5-Sulfamoyl-1,3,4-thiadiazol-2-yl)acetamide and has the following chemical structure:

[structure]

MW 222.24

$C_4H_6N_4O_3S_2$

DIAMOX SEQUELS are extended-release capsules, for oral administration, each containing 500 mg of acetazolamide and the following inactive ingredients:

Microcrystalline cellulose, sodium lauryl sulfate and talc.

The ingredients in the capsule shell are D&C red no. 28, D&C yellow no. 10, FD&C red no. 40, gelatin and titanium dioxide.

The ingredients in the imprinting ink are D&C yellow no. 10 aluminum lake, FD&C blue no. 1 aluminum lake, FD&C blue no. 2 aluminum lake, FD&C red no. 40 aluminum lake, pharmaceutical glaze, propylene glycol and synthetic iron oxide.

**CLINICAL PHARMACOLOGY**

DIAMOX is a potent carbonic anhydrase inhibitor, effective in the control of fluid secretion (e.g., some types of glaucoma), in the treatment of certain convulsive disorders (e.g., epilepsy) and in the promotion of diuresis in instances of abnormal fluid retention (e.g., cardiac edema).

DIAMOX is not a mercurial diuretic. Rather, it is a non-bacteriostatic sulfonamide possessing a chemical structure and pharmacological activity distinctly different from the bacteriostatic sulfonamides.

DIAMOX is an enzyme inhibitor that acts specifically on carbonic anhydrase, the enzyme that catalyzes the reversible reaction involving the hydration of carbon dioxide and the dehydration of carbonic acid. In the eye, this inhibitory action of acetazolamide decreases the secretion of aqueous humor and results in a drop in intraocular pressure, a reaction considered desirable in cases of glaucoma and even in certain non-glaucomatous conditions. Evidence seems to indicate that DIAMOX has utility as an adjuvant in treatment of certain dysfunctions of the central nervous system (e.g., epilepsy). Inhibition of carbonic anhydrase in this area appears to retard abnormal, paroxysmal, excessive discharge from central nervous system neurons. The diuretic effect of DIAMOX is due to its action in the kidney on the reversible reaction involving hydration of carbon dioxide and dehydration of carbonic acid. The result is renal loss of  $HCO_3$  ion, which carries out sodium, water, and potassium. Alkalinization of the urine and promotion of diuresis are thus affected. Alteration in ammonia

metabolism occurs due to increased reabsorption of ammonia by the renal tubules as a result of urinary alkalinization.

DIAMOX SEQUELS provide prolonged action to inhibit aqueous humor secretion for 18 to 24 hours after each dose, whereas tablets act for only eight to 12 hours. The prolonged continuous effect of SEQUELS permits a reduction in dosage frequency.

Plasma concentrations of acetazolamide peak from three to six hours after administration of DIAMOX SEQUELS, compared to one to four hours with tablets. Food does not affect bioavailability of DIAMOX SEQUELS.

Placebo-controlled clinical trials have shown that prophylactic administration of DIAMOX at a dose of 250 mg every eight to 12 hours (or a 500 mg controlled-release capsule once daily) before and during rapid ascent to altitude results in fewer and/or less severe symptoms of acute mountain sickness (AMS) such as headache, nausea, shortness of breath, dizziness, drowsiness, and fatigue. Pulmonary function (e.g., minute ventilation, expired vital capacity, and peak flow) is greater in the DIAMOX treated group, both in subjects with AMS and asymptomatic subjects. The DIAMOX treated climbers also had less difficulty in sleeping.

### **INDICATIONS AND USAGE**

For adjunctive treatment of: chronic simple (open-angle) glaucoma, secondary glaucoma, and preoperatively in acute angle-closure glaucoma where delay of surgery is desired in order to lower intraocular pressure. DIAMOX is also indicated for the prevention or amelioration of symptoms associated with acute mountain sickness despite gradual ascent.

### **CONTRAINDICATIONS**

Hypersensitivity to acetazolamide or any excipients in the formulation. Since acetazolamide is a sulfonamide derivative, cross sensitivity between acetazolamide, sulfonamides and other sulfonamide derivatives is possible.

Acetazolamide therapy is contraindicated in situations in which sodium and/or potassium blood serum levels are depressed, in cases of marked kidney and liver disease or dysfunction, in suprarenal gland failure, and in hyperchloremic acidosis. It is contraindicated in patients with cirrhosis because of the risk of development of hepatic encephalopathy.

Long-term administration of DIAMOX is contraindicated in patients with chronic non-congestive angle-closure glaucoma since it may permit organic closure of the angle to occur while the worsening glaucoma is masked by lowered intraocular pressure.

### **WARNINGS**

Fatalities have occurred, although rarely, due to severe reactions to sulfonamides including Stevens-Johnson syndrome, toxic epidermal necrolysis, fulminant hepatic necrosis, anaphylaxis, agranulocytosis, aplastic anemia, and other blood dyscrasias. Sensitizations may recur when a sulfonamide is readministered irrespective of the route of administration. If signs of hypersensitivity or other serious reactions occur, discontinue use of this drug.

Caution is advised for patients receiving concomitant high-dose aspirin and DIAMOX, as anorexia, tachypnea, lethargy, metabolic acidosis, coma, and death have been reported.

## **PRECAUTIONS**

### **General**

Increasing the dose does not increase the diuresis and may increase the incidence of drowsiness and/or paresthesia. Increasing the dose often results in a decrease in diuresis. Under certain circumstances, however, very large doses have been given in conjunction with other diuretics in order to secure diuresis in complete refractory failure.

### **Information for Patients**

Adverse reactions common to all sulfonamide derivatives may occur: anaphylaxis, fever, rash (including erythema multiforme, Steven-Johnson syndrome, toxic epidermal necrolysis), crystalluria, renal calculus, bone marrow depression, thrombocytopenic purpura, hemolytic anemia, leukopenia, pancytopenia, and agranulocytosis. Caution is advised for early detection of such reactions and the drug should be discontinued and appropriate therapy instituted.

In patients with pulmonary obstruction or emphysema where alveolar ventilation may be impaired, DIAMOX which may precipitate or aggravate acidosis should be used with caution. Gradual ascent is desirable to try to avoid acute mountain sickness. If rapid ascent is undertaken and DIAMOX is used, it should be noted that such use does not obviate the need for prompt descent if severe forms of high altitude sickness occur, i.e., high altitude pulmonary edema (HAPE) or high altitude cerebral edema.

Caution is advised for patients receiving concomitant high-dose aspirin and DIAMOX, as anorexia, tachypnea, lethargy, metabolic acidosis, coma, and death have been reported (see **WARNINGS**).

Both increases and decreases in blood glucose have been described in patients treated with acetazolamide. This should be taken into consideration in patients with impaired glucose tolerance or diabetes mellitus.

Acetazolamide treatment may cause electrolyte imbalances, including hyponatremia and hypokalemia, as well as metabolic acidosis. Therefore, periodic monitoring of serum electrolytes is recommended. Particular caution is recommended in patients with conditions that are associated with, or predispose a patient to, electrolyte and acid/base imbalances, such as patients with impaired renal function (including elderly patients; see **PRECAUTIONS, Geriatric Use**), patients with diabetes mellitus, and patients with impaired alveolar ventilation.

Some adverse reactions to acetazolamide, such as drowsiness, fatigue, and myopia, may impair the ability to drive and operate machinery.

### **Laboratory Tests**

To monitor for hematologic reactions common to all sulfonamides, it is recommended that a baseline CBC and platelet count be obtained on patients prior to initiating DIAMOX therapy and at regular intervals during therapy. If significant changes occur, early discontinuance and institution of appropriate therapy are important. Periodic monitoring of serum electrolytes is recommended.

### **Drug Interactions**

Aspirin- See **WARNINGS**

DIAMOX modifies phenytoin metabolism with increased serum levels of phenytoin. This may increase or enhance the occurrence of osteomalacia in some patients receiving chronic phenytoin therapy. Caution is advised in patients receiving chronic concomitant therapy. By decreasing the

gastrointestinal absorption of primidone, DIAMOX may decrease serum concentrations of primidone and its metabolites, with a consequent possible decrease in anticonvulsant effect. Caution is advised when beginning, discontinuing, or changing the dose of DIAMOX in patients receiving primidone.

Because of possible additive effects with other carbonic anhydrase inhibitors, concomitant use is not advisable.

Acetazolamide may increase the effects of other folic acid antagonists.

Acetazolamide decreases urinary excretion of amphetamine and may enhance the magnitude and duration of their effect.

Acetazolamide reduces urinary excretion of quinidine and may enhance its effect.

Acetazolamide may prevent the urinary antiseptic effect of methenamine.

Acetazolamide increases lithium excretion and the lithium may be decreased.

Acetazolamide and sodium bicarbonate used concurrently increases the risk of renal calculus formation.

Acetazolamide may elevate cyclosporine levels.

#### **Drug/laboratory test interactions**

Sulfonamides may give false negative or decreased values for urinary phenolsulfonphthalein and phenol red elimination values for urinary protein, serum non-protein, and serum uric acid.

Acetazolamide may produce an increased level of crystals in the urine.

Acetazolamide interferes with the HPLC method of assay for theophylline. Interference with the theophylline assay by acetazolamide depends on the solvent used in the extraction; acetazolamide may not interfere with other assay methods for theophylline.

#### **Carcinogenesis, Mutagenesis, Impairment of Fertility**

Long-term studies in animals to evaluate the carcinogenic potential of DIAMOX have not been conducted. In a bacterial mutagenicity assay, DIAMOX was not mutagenic when evaluated with and without metabolic activation.

The drug had no effect on fertility when administered in the diet to male and female rats at a daily intake of up to 4 times the recommended human dose of 1000 mg in a 50 kg individual.

#### **Pregnancy: Teratogenic effects: Pregnancy Category C**

Acetazolamide, administered orally or parenterally, has been shown to be teratogenic (defects of the limbs) in mice, rats, hamsters, and rabbits. There are no adequate and well-controlled studies in pregnant women. Acetazolamide should be used in pregnancy only if the potential benefit justifies the potential risk to the fetus.

#### **Nursing Mothers**

Because of the potential for serious adverse reactions in nursing infants from DIAMOX, a decision should be made whether to discontinue nursing or to discontinue the drug taking into account the

importance of the drug to the mother. Acetazolamide should only be used by nursing women if the potential benefit justifies the potential risk to the child.

### **Pediatric Use**

The safety and effectiveness of DIAMOX SEQUELS in pediatric patients below the age of 12 years have not been established. Growth retardation has been reported in children receiving long-term therapy, believed secondary to chronic acidosis.

### **Geriatric Use**

Metabolic acidosis, which can be severe, may occur in the elderly with reduced renal function.

In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal, or cardiac function, and concomitant disease or other drug therapy.

### **ADVERSE REACTIONS**

**Body as a whole:** Headache, malaise, fatigue, fever, pain at injection site, flushing, growth retardation in children, flaccid paralysis, anaphylaxis.

**Digestive:** Gastrointestinal disturbances such as nausea, vomiting, diarrhea.

**Hematological/Lymphatic:** Blood dyscrasias such as aplastic anemia, agranulocytosis, leukopenia, thrombocytopenic purpura, melena.

**Hepato-biliary disorders:** Abnormal liver function, cholestatic jaundice, hepatic insufficiency, fulminant hepatic necrosis

**Metabolic/Nutritional:** Metabolic acidosis, electrolyte imbalance, including hypokalemia, hyponatremia, osteomalacia with long-term phenytoin therapy, loss of appetite, taste alteration, hyper/hypoglycemia

**Nervous:** Drowsiness, paresthesia (including numbness and tingling of extremities and face), depression, excitement, ataxia, confusion, convulsions dizziness

**Skin:** Allergic skin reactions including urticaria, photosensitivity, Stevens-Johnson syndrome, toxic epidermal necrolysis

**Special senses:** Hearing disturbances, tinnitus, transient myopia

**Urogenital:** Crystalluria, increased risk of nephrolithiasis with long-term therapy, hematuria, glycosuria, renal failure polyuria

### **OVERDOSAGE**

No specific antidote is known. Treatment should be symptomatic and supportive.

Electrolyte imbalance, development of an acidotic state, and central nervous system effects might be expected to occur. Serum electrolyte levels (particularly potassium) and blood pH levels should be monitored.

Supportive measures are required to restore electrolyte and pH balance. The acidotic state can usually be corrected by the administration of bicarbonate.

Despite its high intraerythrocytic distribution and plasma protein binding properties, DIAMOX may be dialyzable. This may be particularly important in the management of DIAMOX overdose when complicated by the presence of renal failure.

## **DOSAGE AND ADMINISTRATION**

### **Glaucoma:**

The recommended dosage is 1 capsule (500 mg) two times a day. Usually 1 capsule is administered in the morning and 1 capsule in the evening. It may be necessary to adjust the dose, but it has usually been found that dosage in excess of 2 capsules (1 g) does not produce an increased effect. The dosage should be adjusted with careful individual attention both to symptomatology and intraocular tension. In all cases, continuous supervision by a physician is advisable.

In those unusual instances where adequate control is not obtained by the twice-a-day administration of DIAMOX SEQUELS, the desired control may be established by means of DIAMOX (tablets or parenteral). Use tablets or parenteral in accordance with the more frequent dosage schedules recommended for these dosage forms, such as 250 mg every four hours, or an initial dose of 500 mg followed by 250 mg or 125 mg every four hours, depending on the case in question.

**Acute Mountain Sickness:** Dosage is 500 mg to 1000 mg daily, in divided doses using tablets or extended-release capsules as appropriate. In circumstances of rapid ascent, such as in rescue or military operations, the higher dose level of 1000 mg is recommended. It is preferable to initiate dosing 24 to 48 hours before ascent and to continue for 48 hours while at high altitude, or longer as necessary to control symptoms.

## **HOW SUPPLIED**

**DIAMOX® SEQUELS®** (Acetazolamide Extended-Release Capsules) are available as 500 mg: Orange opaque cap and orange opaque body filled with white to off-white pellets. Imprinted in black ink, Barr 699. Available in bottles of:  
100

NDC 51285-754-02

Store at controlled room temperature 20° to 25°C (68° to 77°F).

DURAMED PHARMACEUTICALS, INC.

Subsidiary of BarrPharmaceuticals, Inc.

Pomona, New York 10970

Revised NOVEMBER 2004

BR-754

Container Label

2120754023502 1-Pharmco305127 2nd 6 PM 3/07 4 208 156 126 0001 1000000

**NDC 51285-754-02**

**Diamox®  
Sequels®**

**(Acetazolamide Extended-Release  
Capsules)**

**500 mg**

**100 Capsules**

**AMC med Rx only**

**NEW  
APPEARANCE**

Store at 20° to 25° C (68° to 77° F) [See USP Controlled Room Temperature].  
Dispense in well-closed containers.

2120754023502

LOT:  
EXP:

2120754023502 R11-04

DURAMED PHARMACEUTICALS, INC.  
Duramed Pharmaceuticals, Inc.  
Raritan, New York 10970  
2120754023502

