

Prototype "Drug Facts" Label

<i>Drug facts</i>	
<i>Active ingredient (in each inhalation)</i>	<i>Purpose</i>
Epinephrine 0.22 mg.....	Bronchodilator
<i>Uses</i>	
<ul style="list-style-type: none"> ■ temporarily relieves shortness of breath, tightness of chest, and wheezing due to bronchial asthma ■ eases breathing for asthma patients by reducing spasms of bronchial muscles 	
<i>Warning</i>	
For inhalation only	
Do not use	
<ul style="list-style-type: none"> ■ unless a doctor has said you have asthma ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. 	
Ask a doctor before use if you have	
<ul style="list-style-type: none"> ■ heart disease ■ thyroid disease ■ diabetes ■ high blood pressure ■ ever been hospitalized for asthma ■ trouble urinating due to an enlarged prostate gland 	
When using this product	
<ul style="list-style-type: none"> ■ overuse may cause nervousness, rapid heart beat, and heart problems ■ do not continue to use, but seek medical assistance immediately if symptoms are not relieved within 20 minutes or become worse ■ do not puncture or throw into incinerator. Contents under pressure. ■ do not use or store near open flame or heat above 120°F (49°C). May cause bursting. 	
Contains CFC 12, 114, substances which harm public health and environment by destroying ozone in the upper atmosphere	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
<i>Directions</i>	
<ul style="list-style-type: none"> ■ do not use more often or at higher doses unless directed by a doctor ■ supervise children using this product ■ adults and children 4 years and over: start with one inhalation, then wait at least 1 minute. If not relieved, use once more. Do not use again for at least 3 hours. ■ Children under 4 years: ask a doctor 	
<i>Other information</i>	
<ul style="list-style-type: none"> ■ store at room temperature, between 20-25°C (68-77°F) ■ see insert for mouthpiece use and care instructions 	
<i>Inactive ingredients</i> alcohol 34%, ascorbic acid, fluorocarbons (propellant), water	
<i>Questions or comments?</i> Call 1- 8 PRIMATENE or 1-877-462-8363 weekdays 9 AM-5 PM EST	
www.Primatene.com	

The prototype label shown above should be followed in content only. The font sizes for title, headings, subheadings, condensed text and other graphic features are required to be in accordance with provisions of 21 CFR 201.66.

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Charles Ganley
8/30/01 12:14:57 PM