

2 Bi-layer Tablets

**NEW**  
**Mucinex**<sup>600 mg</sup>  
Guaifenesin Extended-Release Tablets  
**EXPECTORANT**

**Drug Facts**

**Active ingredient (in each extended-release tablet) Purpose**  
Guaifenesin 600 mg ..... Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Do not use** ■ for children under 12 years of age

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if** ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Drug Facts (continued)**

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

**Other information**

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NDC 63824-008-82

**NEW**  
**Mucinex**<sup>600 mg</sup>  
Guaifenesin Extended-Release Tablets  
**EXPECTORANT**



2 Bi-layer Tablets



**Mucinex**<sup>600 mg</sup>  
Guaifenesin Extended-Release Tablets  
**EXPECTORANT**




Please visit our Web site:  
[www.adamslabs.com](http://www.adamslabs.com)

PHYSICIAN SAMPLE-  
NOT TO BE SOLD



Fort Worth, Texas 76155 022802  
© 2002 Adams Laboratories, Inc. 2000802

3-63824-008-82-0



NDC 63824-008-82

**Mucinex**<sup>®</sup>  
600mg  
Guaifenesin Extended-Release Tablets

**EXPECTORANT**

2 Bi-layer Tablets

SEE CARTON FOR COMPLETE INFORMATION

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

**PHYSICIAN  
SAMPLE-  
NOT TO BE SOLD**

**Adams**<sup>®</sup>  
LABORATORIES, INC.  
Fort Worth, Texas 76155

3100802 022802

3-6382400882-0

NDC 63824-008-10

**Mucinex**<sup>600mg</sup>  
Guaifenesin Extended-Release Tablets  
**EXPECTORANT**

3-6382400810-3

100 Bi-layer Tablets

**Adams**<sup>®</sup>  
LABORATORIES, INC. Fort Worth, Texas 76155  
3000810 022802

### Drug Facts

**Active ingredient (in each extended-release tablet) Purpose**  
Guaifenesin 600 mg ..... Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

#### Warnings

**Do not use** ■ for children under 12 years of age

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if**

- cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding, ask a health professional before use.** ▶

### Drug Facts (continued)

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

#### Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

#### Other information

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

20 BI-LAYER TABLETS

**Mucinex**  
600mg  
NEW  
Guaifenesin Extended-Release Tablets  
EXPECTORANT

**Drug Facts**

**Active ingredient Purpose**  
(in each extended-release tablet)  
Guaifenesin 600 mg . . . . . Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**  
**Do not use** ■ for children under 12 years of age  
**Ask a doctor before use if you have**  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
■ cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if** ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▶

**NEW**  
**Mucinex**<sup>600mg</sup>  
Guaifenesin Extended-Release Tablets  
EXPECTORANT

**Drug Facts (continued)**  
**Directions**  
■ do not crush, chew, or break tablet  
■ take with a full glass of water  
■ this product can be administered without regard for the timing of meals  
■ adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.  
■ children under 12 years of age: do not use

**Other information**  
■ tamper evident: do not use if neckband is broken  
■ store between 20-25°C (68-77°F)  
■ see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NDC 63824-008-20

**NEW**  
**Mucinex**<sup>600mg</sup>  
Guaifenesin Extended-Release Tablets  
EXPECTORANT



20 BI-LAYER TABLETS

**NEW**  
**Mucinex**<sup>600mg</sup>  
Guaifenesin Extended-Release Tablets  
EXPECTORANT

Please visit our Web site:  
[www.adamslabs.com](http://www.adamslabs.com)

**Adams**  
LABORATORIES, INC.

Fort Worth, Texas 76155  
© 2002 Adams Laboratories, Inc.

030802  
2000820

3-63824-008-20-2

NDC 63824-008-20

**Mucinex**<sup>®</sup> 600mg  
Guaifenesin Extended-Release Tablets  
**EXPECTORANT**



20 Bi-layer Tablets

**Adams**<sup>®</sup>  
LABORATORIES, INC. Fort Worth, Texas 76155  
3000820 030802

**Drug Facts**

**Active ingredient (in each extended-release tablet) Purpose**  
Guaifenesin 600 mg ..... Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Do not use** ■ for children under 12 years of age

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if** ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use. ▶

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

**Other information**

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

40 BI-LAYER TABLETS

**Mucinex**  
600mg  
NEW  
Guaifenesin Extended-Release Tablets  
EXPECTORANT

**Drug Facts**

**Active ingredient Purpose**  
**(in each extended-release tablet)**  
Guaifenesin 600 mg . . . . . Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**  
**Do not use** ■ for children under 12 years of age  
**Ask a doctor before use if you have**  
■ persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema  
■ cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if** ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. ▶

**NEW**  
**Mucinex**  
600mg  
Guaifenesin Extended-Release Tablets  
EXPECTORANT

**Drug Facts (continued)**  
**Directions**  
■ do not crush, chew, or break tablet  
■ take with a full glass of water  
■ this product can be administered without regard for the timing of meals  
■ adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.  
■ children under 12 years of age: do not use

**Other information**  
■ tamper evident: do not use if neckband is broken  
■ store between 20-25°C (68-77°F)  
■ see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

**NEW** NDC 63824-008-40  
**Mucinex**  
600mg  
Guaifenesin Extended-Release Tablets  
EXPECTORANT



40 BI-LAYER TABLETS

**NEW**  
**Mucinex**  
600mg  
Guaifenesin Extended-Release Tablets  
EXPECTORANT

Please visit our Web site:  
[www.adamslabs.com](http://www.adamslabs.com)

**Adams**  
LABORATORIES, INC.  
Fort Worth, Texas 76155  
© 2002 Adams Laboratories, Inc.  
022802  
2000840

3-63824-008-40-0

NDC 63824-008-40

**Mucinex**<sup>®</sup> 600mg  
Guaifenesin Extended-Release Tablets  
**EXPECTORANT**



40 Bi-layer Tablets

**Adams**<sup>®</sup>  
LABORATORIES, INC. Fort Worth, Texas 76155  
3000840 022802

**Drug Facts**

**Active ingredient (in each extended-release tablet) Purpose**  
Guaifenesin 600 mg ..... Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

**Warnings**

**Do not use** ■ for children under 12 years of age

**Ask a doctor before use if you have**

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if** ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding,** ask a health professional before use. ▶

**Drug Facts (continued)**

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

**Other information**

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF

NDC 63824-008-50

# Mucinex<sup>600mg</sup>

Guaifenesin Extended-Release Tablets

**EXPECTORANT**

3-6382400850-9



500 Bi-layer Tablets

**Adams**<sup>®</sup>  
LABORATORIES, INC. Fort Worth, Texas 76155

**Adams**<sup>®</sup>  
LABORATORIES, INC.

3000650 022802

## Drug Facts

**Active ingredient (in each extended-release tablet) Purpose**  
Guaifenesin 600 mg ..... Expectorant

**Uses** helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

### Warnings

**Do not use** ■ for children under 12 years of age

#### Ask a doctor before use if you have

- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema
- cough accompanied by too much phlegm (mucus)

**Stop use and ask a doctor if** ■ cough lasts more than 7 days, comes back, or occurs with fever, rash, or persistent headache. These could be signs of a serious illness.

**If pregnant or breast-feeding**, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

### Directions

- do not crush, chew, or break tablet
- take with a full glass of water
- this product can be administered without regard for the timing of meals
- adults and children 12 years of age and over: one or two tablets every 12 hours. Do not exceed 4 tablets in 24 hours.
- children under 12 years of age: do not use

### Other information

- tamper evident: do not use if neckband is broken
- store between 20-25°C (68-77°F)
- see bottom of bottle for lot code and expiration date

**Inactive ingredients** carbomer 934P, NF; dyes; hydroxypropyl methylcellulose, USP; magnesium stearate, NF; microcrystalline cellulose, NF; sodium starch glycolate, NF



-----  
**This is a representation of an electronic record that was signed electronically and  
this page is the manifestation of the electronic signature.**  
-----

/s/

-----  
Badrul Chowdhury  
7/12/02 03:44:27 PM

Linda Katz  
7/12/02 03:49:12 PM  
Linda M. Katz, M.D. signing for Charles J. Ganley, M.D.