

Drug Facts

Active ingredient (in each applicator) Purpose
 Tioconazole 300 mg (6.5%) Vaginal antifungal

Use • treats vaginal yeast infections

Warnings
 For vaginal use only

Do not use

- if you have never had a vaginal yeast infection diagnosed by a doctor
- if you have a fever (higher than 100 F), pain in the lower abdomen, back, or either shoulder, or foul-smelling vaginal discharge. You should see a doctor for these symptoms.

Ask a doctor before use if you have

- vaginal itching and discomfort for the first time
- symptoms that return within 2 months. You could be pregnant, or have a serious underlying medical cause for your symptoms, including diabetes or a weakened immune system (which may be due to HIV - the virus that causes AIDS).

When using this product

- do not use tampons, douches, spermicides, or other vaginal products
- do not use condoms, or diaphragms, as this product may damage them and they may not prevent sexually transmitted disease or pregnancy.
- do not have vaginal intercourse
- wait 3 days after treatment to resume using condoms or a diaphragm.

Stop use and ask a doctor if

- symptoms do not get better after 3 days
- symptoms last more than 7 days. You may have a condition other than a yeast infection
- you get a rash, fever, abdominal pain, or a foul-smelling vaginal discharge

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- before using, read the enclosed brochure and instructions on foil packet
- adults and children 12 years and over: open the foil packet just before use and remove blue cap. Insert entire contents of applicator into the vagina, preferably at bedtime.
- children under 12: ask a doctor

Other information

- this product is a 1-dose treatment, most women do not experience complete relief of their symptoms in just one day. Most women experience some improvement within 1 day and complete relief of symptoms within 7 days.
- if you have questions about vaginal yeast infections, consult your doctor
- store at 15°-30°C (59°-86°F)
- see left side of foil package for lot number and expiration date

Inactive ingredients butylated hydroxyanisole, magnesium aluminum silicate, white petrolatum

Questions or comments? 1-877-666-4782

Distributed by: Personal Products Company
 Division of McNeil-PPC Inc., Skillman, NJ 08558-9418
 © McNeil-PPC, Inc. 2001
 XX XXXX-XX-XX

Tear Here
 Do not use if sealed 1-DAY foil packet is torn, open or incompletely sealed


1-Dose Treatment

1-DAY


TIOCONAZOLE OINTMENT 6.5%
 VAGINAL ANTIFUNGAL

DIRECTIONS FOR USING 1-DAY

- Tear open foil packet just before using. It is best to use at bedtime.
- Remove applicator and plunger from packet. Applicator is prefilled with 1-DAY vaginal ointment.
- While firmly holding the blue-capped end of the applicator, push the tip of the plunger into the base of applicator.



REMOVE BLUE CAP from top of applicator with a pull-twist action.



Insert 1-DAY

- Lie on your back with your knees bent. Gently insert applicator into the vagina as far as it will go comfortably.
- Push the plunger into the applicator until it will go no farther. Withdraw the applicator and plunger and dispose of it in the wastebasket. Do not flush.
- **DO NOT USE TAMPONS** while using this medicine. Use sanitary napkins instead.

PLEASE READ EDUCATIONAL BROCHURE FOR ADDITIONAL INFORMATION.

**This is a representation of an electronic record that was signed electronically and
this page is the manifestation of the electronic signature.**

/s/

Linda Katz
8/9/01 01:54:01 PM