

Other information ● each caplet contains: sodium 20 mg ● store at 20-25° C (68-77° F). Avoid

**Drug Facts** 

Active ingredient (in each caplet)

Naproxen sodium 220 mg (naproxen 200 mg)....

Drug Facts (continued)

high humidity and excessive heat above 40°C (104°F).

Purpose

Allergy alert: Naproxen sodium may cause a severe allergic reaction which may include: . hives facial swelling ● asthma (wheezing) ● shock

Pain reliever

**Uses** temporarily relieves minor aches and pains due to menstrual cramps muscle aches headache
 backache

## Drug Facts (continued)

Alcohol warning: If you consume 3 or more alcoholic drinks every day, ask your doctor whether you should take naproxen sodium or other pain relievers/fever reducers. Naproxen sodium may cause stomach bleeding.

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if you have ever had serious side effects from any pain reliever

## Ask a doctor or pharmacist before use if you are

- under a doctor's care for any serious condition.
- taking any other drug
   taking any other product that contains naproxen sodium, or any other pain reliever/fever reducer

Stop use and ask a doctor if an allergic reaction occurs. Seek medical help right away.

- pain gets worse or lasts more than 10 days
- you have difficulty swallowing
   it feels like the pill is stuck in your throat . you develop heartburn
- stomach pain occurs or lasts, even if symptoms are mild redness or swelling is present in the painful area 
  any new symptoms appear

## Drug Facts (continued)

Drug Facts (continued)

Warnings

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use naproxen sodium during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

## Directions . do not take more than directed

drink a full glass of water with each dose

under 12 years ask a doctor

12 to 65 years • take 1 caplet every 8 to 12

hours while symptoms last. For the first dose you may take 2 caplets within the first hour. The smallest effective dose should be used. Do not exceed 2 caplets in any 8- to 12-hour period. Do not exceed 3 caplets in a 24-hour period.

over age 65

 do not take more than 1 caplet every 12 hours unless directed by a doctor

MAXIMUM STRENGTH Midal® Extended Relief Dist. by: Bayer HealthCare READ CARTON BEFORE USE. Each caplet contains: Maproxen sodium 220 mg 8875440 MAXIMUM STRENGTH

Midel® Extended Relief

READ CARTON BEFORE USE. Each caplet contains: Magroxen sodium 220 mg 8875440

READ CARTON BEFORE READ CARTON BEFORE
USE. Each caplet contains:
Naprexen sodium 220 mg 12

MAXIMUM STRENGTH Midal@ Extended Relief

HAXIMUM STRENGTH Midal® Extended Relief
Dist. by: Bayer HealthCan Dist. by: Bayer HealthCare READ CARTON BEFORE

USE. Each caplet contains: Naproxen sodium 220 mg 8875440

MAXIMUM STRENGTH Midal@ Extended Relief READ CARTON BEFORE USE. Each caplet contains: See Naproxen sodium 220 mg 3875440

HAXIMUM STRENGTH Midal® Extended Relief READ CARTON BEFORE USE. Each caplet contains: Naproxon sodium 220 mg 8575440